

**NYSCSH Summary of Key Facts 2019-2020 School Year  
 NYSDOH Immunization Requirements for School Entrance/Attendance**

**Prekindergarten (Day Care, Head Start, Nursery or Pre-K)**

Diphtheria and Tetanus toxoid-containing Vaccine and Pertussis Vaccine (DTaP/DTP)	<b>4 doses</b>
Polio Vaccine (IPV/OPV)	<b>3 doses</b>
Measles, Mumps and Rubella Vaccine (MMR)	<b>1 dose</b>
Hepatitis B Vaccine	<b>3 doses</b>
Varicella (Chickenpox) Vaccine	<b>1 dose</b>
Hemophilus influenzae type b conjugate vaccine (Hib)	<b>1 to 4 doses</b>
Pneumococcal Vaccine (PCV)	<b>1 to 4 doses</b>

**Kindergarten through Grade 5**

Diphtheria and Tetanus toxoid-containing Vaccine and Pertussis Vaccine (DTaP/DTP)	<b>5 doses or 4 doses</b> if the 4 <sup>th</sup> dose was received at 4 yrs old or <b>3 doses</b> if 7 yrs old or older and started the series after 1 yr old or older
Polio Vaccine (IPV/OPV)	<b>4 doses or 3 doses</b> if the 3 <sup>rd</sup> dose was received at 4 yrs old or older and at least 6 months from previous dose.
Measles, Mumps and Rubella Vaccine (MMR)	<b>2 doses</b> (2 measles, 2 mumps 1 rubella)
Hepatitis B Vaccine	<b>3 doses</b>
Varicella (Chickenpox) Vaccine	<b>2 doses</b>

**Grades 6 through 11**

Diphtheria and Tetanus toxoid-containing Vaccine and Pertussis Vaccine (DTaP/DTP)	<b>3 doses</b>
Tetanus and Diphtheria toxoid-containing Vaccine and Pertussis Vaccine booster (Tdap)	<b>1 dose</b>
Polio Vaccine (IPV/OPV)	<b>4 doses or 3 doses</b> if the 3 <sup>rd</sup> dose was received at 4 years of age or older
Measles, Mumps and Rubella Vaccine (MMR)	<b>2 doses</b> (2 measles, 2 mumps 1 rubella)
Hepatitis B Vaccine	<b>3 doses or 2 doses</b> of adult hepatitis B vaccine(Recombivax) for children who received the doses at least 4 months apart between the ages of 11 - 15 years of age
Varicella (Chickenpox) Vaccine	<b>2 doses</b>
Meningococcal Vaccine <b>(For Grades 7, 8, 9, &amp; 10 ONLY)</b>	<b>1 dose</b> in grade 7 if not received in grade 7, must obtain in grade 8, 9, and 10 (per move up schedule)

**Grade 12**

Diphtheria and Tetanus toxoid-containing Vaccine and Pertussis Vaccine (DTaP/DTP)	<b>3 doses</b>
Tetanus and Diphtheria toxoid-containing Vaccine and Pertussis Vaccine booster (Tdap)	<b>1 dose</b>
Polio Vaccine (IPV/OPV)	<b>3 doses</b>
Measles, Mumps and Rubella Vaccine (MMR)	<b>2 doses</b> (2 measles, 2 mumps 1 rubella)
Hepatitis B Vaccine	<b>3 doses or 2 doses</b> of adult hepatitis B vaccine (Recombivax) for children who received the doses at least 4 months apart between the ages of 11 - 15 years of age
Varicella (Chickenpox) Vaccine	<b>1 dose</b>
Meningococcal Vaccine	<b>2 doses</b> in grade 12 unless <b>1 dose</b> was given on or after <i>Age 16</i>