






















I NEED TO STAY AT HOME IF....

						
FEVER	VOMITING	DIARRHEA	RASH	HEAD LICE	EYE INFECTION	HOSPITAL STAY AND/OR ER VISIT
Temperature of 100.4 or higher	Within the past 24 hours	Within the past 24 hours	Body rash with itching or fever	Itchy head, active head lice	Redness, itching, and/or "crusty" drainage from eye	Hospital stay and/or ER visit

I AM READY TO GO BACK TO SCHOOL WHEN I AM...

						
FEVER FREE	VOMITING FREE	DIARRHEA FREE	RASH FREE	HEAD LICE FREE	EYE INFECTION FREE	HOSPITAL STAY AND/OR ER VISIT
Fever free for 24 hours without the use of fever reducing medication i.e. Tylenol, Motrin	Free from vomiting for at least 2 solid meals	Free from diarrhea for at least 24 hours	Free from rash itching, or fever. I have been evaluated by my doctor if needed.	Treated with appropriate lice treatment at home and proof is provided to nurse.	Evaluated by my doctor and have note to return to school.	Released by medical provider to return to work or school.

NECESITO QUEDARME EN CASA SI...

TENGO FIEBRE	ESTOY VOMITANDO	TENGO DIARREA	TENGO SARPUILLIDO	TENGO PIOJOS EN LA CABEZA	TENGO UNA INFECCIÓN DE OJO	ESTUVE EN EL HOSPITAL
						
Temperatura de 100° F o más	Dentro de las últimas 24 horas	Dentro de las últimas 24 horas	Sarpullido corporal con fiebre o picazón	Picazón en la cabeza, piojos activos	Enrojecimiento, picazón y/o drenaje de pus del ojo	Estando en hospital y/o visita a sala de urgencias

ESTOY LISTO PARA VOLVER A LA ESCUELA AL SER ...

Libre de fiebre sin la ayuda de medicamentos durante 24 horas (es decir, Tylenol, Motrin, Advil)	Libre de vómitos durante 24 horas.	Libre de diarrea durante 24 horas.	Libre de sarpullido, picazón o fiebre. He sido evaluado por mi médico si es necesario.	Tratado con tratamiento de piojos apropiado en casa.	Libre de drenaje y/o he sido evaluado por mi médico, si fuera necesario.	Dado de alta por mi proveedor de servicios médicos para regresar a la escuela.
--	------------------------------------	------------------------------------	--	--	--	--

Si su hijo tiene faringitis estreptocócica u otra infección bacteriana, debe quedarse en casa hasta haberle administrado el antibiótico durante al menos 24 horas y su proveedor de atención médica le haya dado permiso para que regrese a la escuela. Le recomendamos que busque atención médica cuando su hijo esté enfermo y siga las recomendaciones de su proveedor de atención médica sobre regresar a la escuela y otras actividades.